

FINITE SUMS OF PROJECTIONS IN PURELY INFINITE SIMPLE C*-ALGEBRAS WITH TORSION K_0

VICTOR KAFTAL, P. W. NG, AND SHUANG ZHANG

ABSTRACT. Assume that \mathcal{A} is a purely infinite simple C*-algebra whose K_0 is a torsion group, namely, contains no free element. Then a positive element $a \in \mathcal{A}$ can be written as a finite sum of projections in \mathcal{A} if and only if either a is a projection or $\|a\| > 1$.

1. Introduction

In [5, Theorem 1.1] Fillmore proved that a positive finite-rank operator $a \in B(H)$ can be expressed as finite sums of projections if and only if $\text{Tr}(a) \in \mathbb{N}$ and $\text{Tr}(a) \geq \text{rank}(a)$, where Tr is the standard integer-valued trace.

An immediate natural question is: which infinite-rank operators can also be expressed as finite sums or infinite sums of projections, where the infinite sums of projections are assumed to converge in the strong topology. Motivated by needs in Frame Theory, this question was posed by Dykema, Freeman, Kornelson, Larson, Ordower, and Weber in ([4]), where they also proved a sufficient condition.

Recently, we obtained in [6, Theorem 1.1] the complete characterization: a positive operator $a \in B(H)$ is a (possibly infinite) sum of projections if and only if

$$\text{Tr}(a_+) \geq \text{Tr}(a_-) \quad \text{and if} \quad \text{Tr}(a_+) < \infty \quad \text{then} \quad \text{Tr}(a_+) - \text{Tr}(a_-) \in \mathbb{N} \cup \{0\}.$$

Here $a_+ := (a - I)\chi_{(1, \infty)}(a)$, and $a_- := (I - a)\chi_{[0, 1)}(a)$ where $\chi(a)$ is the spectral measure of a .

A similar characterization, but without the integrality condition, can be given for all positive diagonalizable operators in σ -finite von Neumann factors. In particular, in type III factors a positive operator a can be expressed as a sum of projections if and only if either a is a projection or $\|a\| > 1$.

Turning into another direction, we generalized the $B(H)$ results to bounded module maps on Hilbert C*-modules. In [7] we considered the multiplier algebras of purely infinite and simple C*-algebras, equipped with the strict topology (in analogy of the strong operator topology on $B(H)$) and obtained the following complete characterization ([7, Theorem 1.1]): if \mathcal{A} is a σ -unital, nonunital, purely infinite simple C*-algebra and a be a positive element of $\mathcal{M}(\mathcal{A})$, then a is a sum of projections belonging to \mathcal{A} , with convergence in the strict topology, if and only if one of the following mutually exclusive conditions hold:

- (i) $\|a\|_e > 1$
- (ii) $\|a\|_e = 1$ and $\|a\| > 1$.
- (iii) $a \in \mathcal{M}(\mathcal{A}) \setminus \mathcal{A}$ is a projection.
- (iv) a is the sum of finitely many projections belonging to \mathcal{A} .

Here $\|\cdot\|_e$ denotes the essential norm on $\mathcal{M}(\mathcal{A})$, namely $\|a\|_e = \|\pi(a)\|$, where $\pi : \mathcal{M}(\mathcal{A}) \rightarrow \mathcal{M}(\mathcal{A})/\mathcal{A}$ is the canonical quotient map.

A harder problem is to characterize the case (iv) above, that is, which positive elements in a C*-algebra \mathcal{A} are finite sums of projections in \mathcal{A} . This question is still open even in $B(H)$.

A sufficient condition for $a \in B(H)^+$ to be a finite sum of projections is that $\|a\|_e > 1$. This result was reported in a survey article [13, Theorem 4.12] by Wu quoting unpublished joint work of Choi and Wu in 1988. The proof of this sufficient condition was presented in their recent paper [1, Theorem 2.2].

The special but important case, αI with $\alpha > 1$, follows also from a delicate analysis in [10], [11], and several other papers which characterized the set

$$\Sigma_n := \{\alpha \in \mathbb{R} \mid \alpha I \text{ is the sum of } n \text{ projections}\}$$

A consequence of their analysis yields that $\bigcup_1^\infty \Sigma_n = [1, \infty)$, i.e., αI is a finite sum of projections if and only if $\alpha \geq 1$.

An independent and different proof of the sufficient condition by Choi and Wu was given by the authors of the present paper in [8]: If M is an infinite σ -finite von Neumann factor and $a \in M$ is positive, then a sufficient condition for a to be a finite sum of projections in M is, respectively,

- (i) $\|a\|_e > 1$ when M is of type I_∞ (where $\|\cdot\|_e$ denotes the usual essential norm of $B(H)$);
- (ii) $\|a\|_e > 1$ when M is of type II_∞ (where $\|\cdot\|_e$ denotes the essential norm relative to the closed ideal $J(M)$ generated by all finite projections of M);
- (iii) $\|a\| > 1$ when M is of type III. Moreover, $a \in M^+$ is a finite sum of projections if and only if either a is a projection or $\|a\| > 1$.

By employing a suitable notion of essential central spectrum, we proved also that a similar sufficient condition holds also for global properly infinite von Neumann algebras [8, Theorem 4.3].

For multiplier algebras we proved in [9, Theorem 3.3] a similar result: if \mathcal{A} is a σ -unital, nonunital, purely infinite simple C^* -algebra, then every positive element $a \in \mathcal{M}(\mathcal{A}) \setminus \mathcal{A}$ with $\|a\|_e > 1$ can be expressed as a finite sum of projections.

Since σ -unital purely infinite simple C^* -algebras share some properties of σ -finite type III factors, one would hope that every positive element a in a purely infinite simple C^* -algebra with $\|a\| > 1$ could also be written as a finite sum of projections. And if not, one would like to see which such algebras do not have this property.

The objective of this paper is to prove the following result.

Theorem 1.1. *Let \mathcal{A} be a purely infinite simple C^* -algebra such that $K_0(\mathcal{A})$ is a torsion group, namely, contains no free elements, then a positive element $a \in \mathcal{A}^+$ is a finite sum of projections if and only if either a is a projection or $\|a\| > 1$.*

Both in the case of von Neumann algebras ([8]) and in the case of multiplier algebras ([7] and [9]), the results were obtained by first decomposing a into a special infinite sum of projections (converging in the strong topology in the von Neumann algebra case, and in the strict topology in the multiplier algebra case), and then, reassembling these sums into finite sums of projections. However, for purely infinite simple C^* -algebras the above techniques can no longer be applied, and we will have to use a different approach.

A key reduction is that if a purely infinite simple C^* -algebra \mathcal{A} has the property that αp is a finite sum of projections for all $\alpha > 1$ and all projections $p \in \mathcal{A}$, then each $a \in \mathcal{A}^+$ with $\|a\| > 1$ is also of a finite sum of projections. The key in this reduction is the fact derived in ([9]) that each positive element $a \in \mathcal{A}$ can be written as positive linear combination of projections.

Then we prove that in any purely infinite simple C^* -algebra, αp is a finite sum of projections for any $\alpha > 1$ and any projection p that represents the zero element of $K_0(\mathcal{A})$ (Proposition 2.3).

This will be done by showing that αp is of a finite sum of projections for any $\alpha \in [\frac{3}{2}, 3]$ (Lemma 2.1), and then, prove that $\alpha p \oplus \beta q$ is a finite sum of projections for any pair of rational numbers $\alpha > 1$ and $\beta \geq 0$ and every projection $p \sim q$, which represents the zero element of $K_0(\mathcal{A})$ (Lemma 2.2).

Combining these results, we then obtain that $\alpha p \oplus \beta q$ is a finite sum of projections for any $\alpha > 1$, $\beta \geq 0$, and any projection $p, q \in \mathcal{A}$ which represent elements of $K_0(\mathcal{A})$ with finite orders (Proposition 2.5).

As a consequence, if all elements of $K_0(\mathcal{A})$ are of finite order, i.e., $K_0(\mathcal{A})$ is a torsion group, then $a \in \mathcal{A}^+$ is a finite sum of projections if and only if either a is a projection or $\|a\| > 1$. (Theorem 1.1).

For the readers' convenience we summarize some basic facts on K -theory of infinite simple C^* -algebras that we will use in this paper. A handy reference are the original papers by Cuntz [2] and [3].

Recall that $[p]$ denotes the equivalence class of a projection $p \in \mathcal{A}$ for any C^* -algebras \mathcal{A} . If \mathcal{A} is infinite and simple, then the K_0 group of \mathcal{A} can be described as follows:

- (1) $K_0(\mathcal{A}) = \{[p] \mid p \text{ is an infinite projection in } \mathcal{A}\}$.
- (2) The addition '+' on $K_0(\mathcal{A})$ is defined as follows: for any two infinite projections $p, q \in \mathcal{A}$, choose $p' \sim p$, $q' \sim q$ so that $p'q' = 0$. then $[p] + [q] = [p' + q']$. In particular, we denote

$$n[p] = \overbrace{[p] + [p] + \cdots + [p]}^{n \text{ times}}.$$

- (3) The zero (or unit) element $\theta \in K_0(\mathcal{A})$ is defined as $[p - p']$ for any pair of projections $p' < p$ with $p' \sim p$ and $p - p'$ infinite.

- (4) For any infinite projection $q \in \mathcal{A}$ and any positive integer m , q can be decomposed into the sum $q = \sum_{j=0}^m q_j$ of mutually orthogonal projections q_j , with $q_0 \sim q$ and $[q_j] = \theta$ for all $j > 0$.
- (5) An infinite C^* -algebra \mathcal{A} has torsion K_0 if for every projections $p \neq 0$ the element $[p]$ is finitely cyclic in $K_0(\mathcal{A})$, namely, there is a positive integer n for which $n[p] = \theta$, the zero element of $K_0(\mathcal{A})$. Among algebras with torsion K_0 are the Cuntz algebras \mathcal{O}_n for $2 \leq n < \infty$, for which $K_0(\mathcal{O}_n) = \mathbb{Z}/(n-1)\mathbb{Z}$; in particular, $K_0(\mathcal{O}_2) = \{\theta\}$, the trivial group.

2. Finite sums of projections and K-theory

The whole section will be devoted to establish the proof of our main result by a sequence of technical lemmas.

Lemma 2.1. *Let \mathcal{A} be an infinite simple C^* -algebra, $p \in \mathcal{A}$ be a projection that represents the zero element of $K_0(\mathcal{A})$ (i.e., $[p] = \theta$) and $\gamma \in [\frac{3}{2}, 3]$. Then there are six projections $q_j \in \mathcal{A}$, with $q_j \sim p$ and such that $\gamma p = \sum_1^6 q_j$.*

Proof. Decompose $p = p_1 + p_2$ into the sum of two mutually orthogonal projections $p_1 \sim p_2 \sim p$. Let

$$\begin{aligned} a_1 &:= (2\gamma - 3)p_1 + (3 - \gamma)p_2 \\ a_2 &:= (3 - \gamma)p_1 + (2\gamma - 3)p_2 \end{aligned}$$

Then

$$\gamma p = a_1 + a_2 \quad \text{and} \quad a_1 \geq 0, \quad a_2 \geq 0.$$

Further decompose $p_2 = p'_2 + p''_2$ into the sum of two mutually orthogonal projections $p'_2 \sim p''_2 \sim p_2$. Then

$$a_1 = (2\gamma - 3)p_1 + (3 - \gamma)p'_2 + (3 - \gamma)p''_2.$$

Since the three projections p_1, p'_2 , and p''_2 are mutually orthogonal and equivalent, a_1 belongs to an embedding of $\mathbb{M}_3(\mathbb{C})$ in $p\mathcal{A}p$ where p_1, p'_2 , and p''_2 are identified with rank-one projections. Thus a_1 can be identified with a positive matrix in $\mathbb{M}_3(\mathbb{C})$ with $\text{Tr}(a_1) = 3$ and $\text{rank}(a_1) \leq 3$. Then it follows from [5, Theorem 1.1] that a_1 is the sum of three projections (all equivalent to p .) The same argument applies to a_2 , and hence, the proof is completed. \square

Lemma 2.2. *Let \mathcal{A} be an infinite simple C^* -algebra, $p, q \in \mathcal{A}$ be two orthogonal projections that represent the zero element of $K_0(\mathcal{A})$, and let α, β be rational numbers with $\alpha > 1$ and $\beta \geq 0$. Then $a := \alpha p + \beta q$ is a finite sum of projections.*

Proof. Set $\alpha = \frac{k}{m}$ and $\beta = \frac{h}{m}$ with k, h, m integers and choose an integer $r \geq \frac{m}{k-m}$, equivalently, $rk \geq rm + m$. Then decompose $p = \sum_1^{rm} p_j$ (resp., $q = \sum_1^m q_j$) into the sum of rm (resp. m) mutually orthogonal projections $p_j \sim q_j \sim p$. Then

$$a = \frac{k}{m} \sum_1^{rm} p_j + \frac{h}{m} \sum_1^m q_j$$

can be identified with a positive matrix in $\mathbb{M}_{rm+m}(\mathbb{C})_+$ with

$$\text{Tr}(a) = rk + h \in \mathbb{N} \quad \text{and} \quad \text{rank}(a) \leq rm + m.$$

Since $rk + h \geq rm + m$, it follows once again from [5, Theorem 1.1] that a is a finite sum of projections. \square

Proposition 2.3. *Let \mathcal{A} be an infinite simple C^* -algebra, $p, q \in \mathcal{A}$ be two orthogonal equivalent projections that represent the zero element of $K_0(\mathcal{A})$, and let α, β be real numbers with $\alpha > 1$ and $\beta \geq 0$. Then $a := \alpha p + \beta q$ is a finite sum of projections.*

Proof. If $\beta = 0$, write $\alpha p = \alpha p_1 + \alpha p_2$, where $p_1 \sim p_2 \sim p$ and $p_1 p_2 = 0$. Thus assume without loss of generality that $\beta > 0$.

Let ρ_1, ρ_2, ρ_3 be rational numbers with

$$\begin{aligned} 1 &< \rho_1 < \alpha \\ \frac{1}{3}(\alpha - \rho_1) &< \rho_2 < \frac{2}{3}(\alpha - \rho_1) \\ \frac{1}{3}\beta &< \rho_3 < \frac{2}{3}\beta. \end{aligned}$$

Decompose $p = p' + p''$ into the sum of two mutually orthogonal projections $p' \sim p'' \sim p$. Then

$$a = \rho_1 p + (\alpha - \rho_1)p'' + (\alpha - \rho_1)p' + \beta q.$$

Since $\frac{\alpha - \rho_1}{\rho_2} \in [\frac{3}{2}, 3]$ and $\frac{\beta}{\rho_3} \in [\frac{3}{2}, 3]$, by Lemma 2.1 we can find projections $\{q_j\}_1^{18}$ with $q_j \sim p$ for all j and such that

$$\begin{aligned} \frac{\alpha - \rho_1}{\rho_2} p'' &= \sum_{j=1}^6 q_j, \\ \frac{\alpha - \rho_1}{\rho_2} p' &= \sum_{j=7}^{12} q_j, \\ \frac{\beta}{\rho_3} q &= \sum_{j=13}^{18} q_j. \end{aligned}$$

It follows that

$$a = \rho_1 p + \rho_2 \sum_{j=1}^6 q_j + \rho_2 \sum_{j=7}^{12} q_j + \rho_3 \sum_{j=13}^{18} q_j$$

Now decompose $p' = \sum_{j=1}^6 p_j$ and $p'' = \sum_{j=7}^{18} p_j$ into the sum of mutually orthogonal projections $p_j \sim p$. Thus

$$a = \sum_{j=1}^6 (\rho_1 p_j + \rho_2 q_j) + \sum_{j=7}^{12} (\rho_1 p_j + \rho_2 q_j) + \sum_{j=13}^{18} (\rho_1 p_j + \rho_3 q_j).$$

Since all the coefficients ρ_i are rational, $\rho_1 > 1$, and $p_j q_j = 0$ for $1 \leq j \leq 18$ because $p' p'' = p q = 0$, it follows by Lemma 2.2 that each of the above 18 summands is a finite sum of projections, and hence, so is a . \square

Taking $\beta = 0$ in the above proposition we thus have:

Corollary 2.4. *Let \mathcal{A} be an infinite simple C^* -algebra and p be a projection in \mathcal{A} that represents the zero element of $K_0(\mathcal{A})$. Then for all $\alpha > 1$ the positive element αp is a finite sum of projections.*

This result extends also to projections that represent finitely cyclic elements of $K_0(\mathcal{A})$.

Proposition 2.5. *Let \mathcal{A} be a purely infinite simple C^* -algebra and p be a projection of \mathcal{A} that represents a finite order element of $K_0(\mathcal{A})$ (i.e., such that $[p]$ is finitely cyclic). Then αp is a finite sum of projections.*

Proof. Assume also without loss of generality that $\alpha < 2$. Let θ be the zero element of $K_0(\mathcal{A})$. By hypothesis $n[p] = \theta$ for some integer $n \in \mathbb{N}$. Then $\frac{1}{\alpha-1} - \frac{1}{n} > 0$ and let $m \geq 0$ be the integer part of $\frac{1}{\alpha-1} - \frac{1}{n}$ and $0 \leq \delta < 1$ be its fractional part, i.e.,

$$m = \frac{1}{\alpha-1} - \frac{1}{n} - \delta.$$

Since $[p] = (m+1)\theta + [p] = ((m+1)n+1)[p]$, we can decompose p into the sum $p = \sum_{j=0}^{(m+1)(n)} p_j$ of mutually orthogonal projection $p_j \sim p$. Then

$$\alpha p = \alpha \sum_{j=0}^{mn} p_j + (\alpha - 1)\delta \sum_{j=mn+1}^{(m+1)n} p_j + (\alpha - (\alpha - 1)\delta) \sum_{j=mn+1}^{(m+1)n} p_j.$$

Let

$$b := \alpha \sum_{j=0}^{mn} p_j + (\alpha - 1)\delta \sum_{j=mn+1}^{(m+1)n} p_j$$

and

$$c := (\alpha - (\alpha - 1)\delta) \sum_{j=mn+1}^{(m+1)n} p_j.$$

Then $a = b + c$. Since all the projections p_j are mutually orthogonal and equivalent (to p), the positive operator b can be identified with a finite matrix with $\text{rank}(b) \leq (m+1)n + 1$ and with

$$\text{Tr}(b) = (mn+1)\alpha + n(\alpha-1)\delta = (m+1)n + 1.$$

Thus, by [5, Theorem 1.1], b is a finite sum of projections. Since $\alpha - (\alpha - 1)\delta > 1$ and since

$$\left[\sum_{j=mn+1}^{(m+1)n} p_j \right] = n[p] = \theta,$$

by Corollary 2.4 c too is a finite sum of projections, and hence, so is a . \square

In particular, if \mathcal{A} is a purely infinite simple C^* -algebra such that $K_0(\mathcal{A})$ is a torsion group, i.e., every element is of finite order, then \mathcal{A} has the property that the element αp is a finite sum of projections for every $\alpha > 1$ and every projection $p \in \mathcal{A}$. As we will see, this property extends to all positive operators with norm larger than 1.

Lemma 2.6. *Let \mathcal{A} be a purely infinite simple C^* -algebra with the property that the element αp is a finite sum of projections for every $\alpha > 1$ and every projection $p \in \mathcal{A}$. Then for every orthogonal pair of projections $0 \neq p, q \in \mathcal{A}$ and every pair of real numbers α, β with $\alpha > 1$ and $\beta \geq 0$, the operator $a = \alpha p + \beta q$ is a finite sum of projections.*

Proof. Without loss of generality assume that $\beta < 1$. Let $n \in \mathbb{N}$ be the integer part of $\frac{2-\beta}{\alpha-1}$ and $0 \leq \delta < 1$ be its fractional part and let $\epsilon := \delta(\alpha - 1)$. Then $0 \leq \epsilon < \alpha - 1$ and

$$n\alpha + \beta + \epsilon = n + 2.$$

Since $[p] = (n+1)[q] + ([p] - (n+1)[q])$ in $K_0(\mathcal{A})$, we can decompose p into the sum $p = p' + \sum_{j=0}^n q_j$ of mutually orthogonal nonzero projections with $q_j \sim q$ for $0 \leq j \leq n$ and a nonzero “remainder” p' also orthogonal to all the projections q_j . Then

$$a = \alpha p' + (\alpha - \epsilon)q_0 + \epsilon q_0 + \sum_{j=1}^n \alpha q_j + \beta q$$

Let $b := \epsilon q_0 + \sum_{j=1}^n \alpha q_j + \beta q$. Then $b \geq 0$ and since all the projections are mutually orthogonal and equivalent, we identify it with a positive matrix with

$$\text{Tr}(b) = n\alpha + \beta + \epsilon = n + 2 \geq \text{rank}(b).$$

Thus b is a finite sum of projections. By hypothesis so are $\alpha p'$ and $(\alpha - \epsilon)q_0$ and hence so is a . \square

Theorem 2.7. *Let \mathcal{A} be a purely infinite simple C^* -algebra with the property that the element αp is a finite sum of projections for every $\alpha > 1$ and every projection $p \in \mathcal{A}$. Then $a \in \mathcal{A}^+$ is a finite sum of projections if and only if either a is a projection or $\|a\| > 1$.*

Proof. If a is a sum of projections, then $\|a\| \geq 1$. Then either the projections are mutually orthogonal, in which case a itself is a projection, or $\|a\| > 1$.

For the converse implication, assume that $\|a\| > 1$. By passing if necessary to the hereditary algebra $(a\mathcal{A}a)^-$ generated by a which is σ -unital, we assume without loss of generality that \mathcal{A} itself is σ -unital. By [9, Remark 2.1 (II)], \mathcal{A} has the PLP property, namely the positive cone of each of its hereditary C^* -subalgebras is the closure of the positive combinations of its projections. But then, by [9, Lemma 2.10], either $a = \alpha p \oplus c$ for some $\alpha > 1$ and nonzero projections $p \in \mathcal{A}$ and a positive element $c \in \mathcal{A}$, or a is the sum of two such operators. Assume without loss of generality that $a = \alpha p \oplus c$. By [9, Theorem 2.11],

$c = \sum_1^n \lambda_j q_j$ for some scalars $\lambda_j > 0$ and projections q_j . Decompose p into the sum $p = \sum_1^n p_j$ of mutually orthogonal nonzero projections $p_j \in \mathcal{A}$. Then

$$a = \sum_1^n (\alpha p_j + \lambda_j q_j)$$

where $p_j q_j = 0$ for all j . By Lemma 2.6, each summand $\alpha p_j + \lambda_j q_j$ is a finite sum of projections, and hence, so is a . □

Combining Proposition 2.5 and Theorem 2.7 we thus obtain our main result.

Theorem 1.1 *Let \mathcal{A} be a purely infinite simple C^* -algebra such that $K_0(\mathcal{A})$ is a torsion group, namely, contains no free elements, then a positive element $a \in \mathcal{A}^+$ is a finite sum of projections if and only if either a is a projection or $\|a\| > 1$.*

Remark 2.8.

- (i) *Purely infinite simple C^* -algebras with $K_0(\mathcal{A})$ a torsion group include of course all those trivial K_0 and all Cuntz algebras \mathcal{O}_n with $2 \leq n < \infty$.*
- (ii) *The proofs of Lemmas 2.1, 2.2, and Proposition 2.3, and hence, the conclusion in Corollary 2.4, hold also in σ -finite von Neumann factors, if restricted to infinite projections. Namely, if M is σ -finite von Neumann factor, then αp is a finite sum of projections for all $\alpha > 1$ and all infinite projections p . This observation provides a simple alternative proof for the fact that $\alpha 1$ is a finite sum of projections in $B(H)$ if $\alpha > 1$ (see [10], [11], [13], [1], [8].)*
- (iii) *Notice explicitly that the techniques of the present paper do not provide an alternative proof for the sufficiency of the condition $\|a\|_e > 1$ for $a \in B(H)^+$ to be a finite sum of projections. Indeed, while by a proof similar to the one for Theorem 1.1 (see [8, Theorem 4.3]) it is enough to consider elements of the form $a = \alpha p \oplus \beta q$, where $\alpha > 1$, $\beta \geq 0$, p, q are projections, and p is infinite, it is not enough to consider only elements where also q is infinite. But the proofs of Proposition 2.3 or of Lemma 2.6 do not apply unless q too is infinite; a different technique is required when q is finite (see [8, Lemma 4.2]).*

Notice also that the condition $\|a\|_e > 1$ implies that $\pi(a)$ is a finite sum of projections in the Calkin algebra $B(H)/K(H)$, but this alone does not imply that the same holds for a (e.g., there are cases of operators $1 + k$ with $k \in K(H)^+$ that are not finite sum of projections (see [8, Example 5.9]).

- (iv) *Technical difficulties occur at attempting the analysis of purely infinite simple C^* -algebras whose K_0 group contains free elements; in particular, even \mathcal{O}_∞ resists all efforts so far. Unlike all type III von Neumann factors in which all hereditary W^* -subalgebras are unital, hereditary C^* -subalgebras of a σ -unital purely infinite simple C^* -algebra \mathcal{A} are either unital or stable (namely, $\mathcal{A} \cong \mathcal{A} \otimes K(H)$), where \mathcal{A} can be chosen unital and $K(H)$ is the stable algebra of all compact operators on a separable Hilbert space (see [14] and [15]). It is well known that infinite rank positive operators in $K(H)$ are not finite sums of projections, but the same positive elements in the canonical embedding of $K(H)$ as a C^* -subalgebra of $\mathcal{A} \otimes K(H)$, become finite sums of projections as long as $K_0(\mathcal{A})$ is a torsion group. Whether the same conclusion can be expected or not, when $K_0(\mathcal{A})$ is not a torsion group, remains mysterious. If not, these purely infinite simple C^* -algebras would exhibit quite a different behavior from type III factors concerning the decompositions of its positive elements.*

Acknowledgement: S. Zhang was supported by a Taft Center Travel Grant when the article was presented in Beijing, China in the summer of 2010.

REFERENCES

- [1] Choi, M.D. and Wu, P.Y., *Sums of orthogonal projections*. Preprint.
- [2] Cuntz, J *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. 57 (1977), 173–185.
- [3] Cuntz, J *K -theory for certain C^* -algebras*, Ann. of Math. 131(1981), 181–197.
- [4] Dykema, K., Freeman, D., Kornelson, K., Larson, D., Ordower, M., and Weber, E., *Ellipsoidal tight frames and projection decompositions of operators*. Illinois J of Math., **48**, (2004), 477–489.
- [5] Fillmore, P., *On sums of projections*. J. Func. Anal., **4** (1969), 146–152.
- [6] Kaftal, V., Ng, P. W., and Zhang, S., *Strong sums of projections in von Neumann factors*. J. Funct. Anal., **257**(2009), 2497–2529.
- [7] Kaftal, V., Ng, P. W., and Zhang, S., *Projection decomposition in multiplier algebras*. Math. Ann., to appear.

- [8] Kaftal, V., Ng, P. W., and Zhang, S., *Finite sums of projections in von Neumann algebras* 2010; preprint
- [9] Kaftal, V., Ng, P. W., and Zhang, S., *Positive combinations and sums of projections in purely infinite simple C^* -algebras and their multiplier algebras*. Preprint.
- [10] Kruglyak, S., Rabanovich V., and Samoilenko, Y., *On Sums of Projections*. Funct Anal and Appl., **36** (2002), 182-195.
- [11] Kruglyak, S., Rabanovich V., and Samoilenko, Y., *Decomposition of a scalar matrix into a sum of orthogonal projections*. Lin. Alg and Applic., **370** (2003), 217-225.
- [12] Marcoux, L., *Projections, commutators and Lie ideals in C^* -algebras* Mathematical Proceedings of the Royal Irish Academy, **110A** (2010), no. 1, 31-55.
- [13] Wu, P.Y., *Additive combinations of special operators*. Funct. Anal Oper. Theory., Banach Ctr Publ, Inst Math Polish Acad Sci., **30** (1994), 337-361.
- [14] Zhang, S., *A property of purely infinite simple C^* -algebras*. Proc. Amer. Math. Soc., **109** (1990) no. 3, 717-720.
- [15] Zhang, S., *Certain C^* -algebras with real rank zero and their corona and multiplier algebras. II* *K-theory* **6** (1992) no. 1, 1-27.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CINCINNATI, P. O. BOX 210025, CINCINNATI, OH, 45221-0025, USA
E-mail address: `kaftalv@ucmail.uc.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LOUISIANA, 217 MAXIM D. DOUCET HALL, P.O. BOX 41010, LAFAYETTE, LOUISIANA, 70504-1010, USA
E-mail address: `png@louisiana.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CINCINNATI, P.O. BOX 210025, CINCINNATI, OH, 45221-0025, USA
E-mail address: `zhangs@math.uc.edu`